



SECOND AND THIRD HARMONIC GENERATION IN METALLIC NANOHYBRIDS

ROBERT GERSTNER¹, MAHI SINGH¹

¹University of Western Ontario, London, Canada

METALLIC NANOPARTICLES





I would like to thank Dr. Singh for his mentorship and contributions to the project, which includes initially deriving the output intensity equations. I would also like to thank his graduate students and particularly Qingzhou Meng for his helpful discussions. I acknowledge our collaborating group from Vanderbilt University who conducted experiments on this effect. Finally, I thank the Western physics and integrated science programs for



Figure 4: The three-layer nanohybrid used in this project.

• Theory fits experiment best for full nanohybrid, worst for

Largest contribution comes from CuS nanolayer, smallest

• CuS offers the most enhancement effects for third har-

¹T. H. Maiman, "Stimulated Optical Radiation in Ruby", Na-

²P. A. Franken, A. E. Hill, C. W. Peters, and G. Weinreich, "Generation of Optical Harmonics", Phys. Rev. Lett. 7, Publisher: American Physical Society, 118–119 (1961).

³R. W. Terhune, P. D. Maker, and C. M. Savage, "Optical Harmonic Generation in Calcite", Phys. Rev. Lett. 8, Publisher:

⁴G. He, Nonlinear optics and photonics, First edition, OCLC: ocn881438178 (Oxford University Press, Oxford, United

⁵L. Mostaço-Guidolin, N. Rosin, and T.-L. Hackett, "Imaging" Collagen in Scar Tissue: Developments in Second Harmonic Generation Microscopy for Biomedical Applications", International Journal of Molecular Sciences 18, 1772 (2017).

⁶H. S. Kim and D. Y. Lee, "Near-Infrared-Responsive Cancer Photothermal and Photodynamic Therapy Using Gold Nanoparticles", Polymers 10, 10.3390/polym10090961

⁷G. Das, M. Coluccio, S. Alrasheed, A. Giugni, M. Allione, B. Torre, G. Perozziello, P. Candeloro, and E. Di Fabrizio, "Plasmonic nanostructures for the ultrasensitive detection